

Graphene mechanics: defects, buckling and domain growth

Sandeep Kumar Jain

Vladimir Juricic and Gerard T. Barkema

Institute for theoretical Physics, Utrecht University, Princetonplein 5, 3584 CC Utrecht, The Netherlands

S.K.Jain@uu.nl

We have developed a new semi-empirical potential for graphene [1], using DFT calculations for determining the various parameters, which for the first time includes a term for out-of-plane deformations. We have demonstrated the usefulness of this potential in studies of different kind of intrinsic defects (Stone-Wales defect, separating dislocations and grain boundaries). Our simulations show that the stress caused by these defects can be relieved by buckling, which extends to hundreds of nanometers. A detailed study of the formation energies of defects surprisingly revealed that the value for the formation energy depends on the type of boundary conditions [2]. Therefore it is necessary to specify the boundary conditions for the energy of the lattice defects in the buckled two-dimensional crystals to be uniquely defined. We have also theoretically described that the vibrational density of states (VDOS) can be used in probing the crystallinity of graphene samples [3]. The novel potential can be effectively combined with interlayer interaction, allowing the simulation of bilayer graphene and study the effect of twist angle on the structure and buckling [4]. Recently, we have described the universal shape behaviour of a graphene gas bubble irrespective of its size. We show

that for small gas bubbles (~10 nm), the vdW pressure is in the order of 1 GPa [5].

References

- [1] S.K.Jain, et. al, JPCC, 199 (2015) 9646
- [2] S.K.Jain, et. al, PRB, 94 (2016) 020102(R)
- [3] S.K.Jain, et. al, JPCL, 6 (2015) 3897
- [4] S.K.Jain, et. al, 2D Mat, 4 (2016) 015018
- [5] S.K.Jain, et. al, Submitted (2017)

Figures




Figure 1: Buckled graphene with a Stone-Wales defect.




Figure 2: (a) Buckling in twisted bilayer graphene. (b) Energy distribution in twisted bilayer graphene and formation of vortices.