

Quantum transport simulations of graphene non-local spin valves

Marc Vila^{1,2}

Christoph W. Groth³, Aron W. Cummings¹,
Stephen R. Power^{1,2}, Xavier Waintal³, Stephan
Roche^{1,4}

¹Catalan Institute of Nanoscience and
Nanotechnology (ICN2), CSIC and The
Barcelona Institute of Science and Technology,
Campus UAB, Bellaterra, 08193 Barcelona, Spain

²Universitat Autònoma de Barcelona, Campus
UAB, Bellaterra, 0819 Barcelona, Spain

³CEA-INAC/UJF Grenoble 1, SPSMS UMR-E 9001,
Grenoble 38054, France

⁴ICREA, Institució Catalana de Recerca i Estudis
Avançats, 08070 Barcelona, Spain

marc.vila@icn2.cat

Graphene is a promising platform for transporting spin information due to its high carrier mobility and low intrinsic spin-orbit coupling. State-of-the-art devices have already demonstrated room temperature long spin lifetimes of several nanoseconds and spin diffusion lengths up to a few tens of μm [1]. The spin relaxation mechanisms have been founded to be strongly monitored by spin-pseudospin entanglement and dephasing effects [2]. The next challenge is to be able to manipulate and control spin signals to engineer active spintronics and hence envision complex graphene-based, non-charged-based devices [3]. Placing graphene on top of ferromagnetic insulators or transition metal dichalcogenides (TMDC) is a promising route to induce, by proximity effect, magnetism or strong spin-orbit coupling (SOC) in graphene while conserving the excellent electronic properties.

In this work, common experimental devices to measure spin-related properties, such as Hanle non-local spin valves (Figure1)[4], are

simulated in graphene within the Landauer-Büttiker formalism using the Kwant python package [5]. Such software is capable of simulating Hanle setups using fully coherent quantum transport. Comparison and discussion with available experimental data are presented for different graphene systems as well as computing local properties like (spin) currents and local density of states. In this way, a more microscopic picture of the spin-dynamics in graphene is obtained [6].

References

- [1] M Drögeler et al., Nano Lett., 16 (2016) 3533
- [2] D Van Tuan et al., Nature Phys., 10 (2014) 857-863; AW Cummings, S Roche, Phys. Rev. Lett., 116 (2016) 086602
- [3] S Roche et al., 2D Mater., 2 (2015) 030202
- [4] B Raes et al., Nat. Comm., 7 (2016) 11444
- [5] CW Groth, M Wimmer, AR Akhmerov and X Waintal, New J. Phys., 16 (2014) 063065
- [6] M. Vila et al, in preparation

Figures




Figure 1: Schematics of the spin precession in a non-local Hanle measurement. Figure extracted from ref. 4.