

Enhancement of local Spin-Orbit Coupling effects in TMDC's/graphene heterostructures mediated by disorder

Armando Pezo

Jose H. Garcia

Adalberto Fazzio

Stephan Roche

Autonomous University of Barcelona,

Edifici ICN2 Campus de la UAB,

Av. de Serragalliners, s/n, 08193 Bellaterra,

Barcelona, Spain

armando.pezo@icn2.cat

On recent years, there is considerable interest in understanding the electronic and transport properties of graphene-based heterostructures. In particular, the heterostructure formed by graphene and two-dimensional transition metal dichalcogenides (TMDC's) is currently under extensive research [1] due to recent measurements of the Spin Hall Effect [2], and the discovery of a giant spin lifetime anisotropy [3]. However, the effects of disorder in these heterostructures is still lacking, mostly due to the high dimensionality of the systems arising from the two incommensurate lattices conforming it. In this work, we use the very recent implementation of the spin-orbit coupling of the SIESTA package [4] deployed in a high-performance computing environment, for showing the effect of disorder in the spin-texture of graphene/TMDC's heterostructures. As our main result, we found that localized defects produced a local modification of the spin-orbit coupling which can be used to increase and tune spin-orbit related phenomena as the Spin Hall Effect.

References

- [1] Jose H. Garcia *et al*, *Chem. Soc. Rev.*, 2018,47, pp 3359-3379.
- [2] C. J. Safeer *et al*, *Nano Lett.*, 2019, 19 (2), pp 1074-1082.
- [3] Talieh S. Ghiasi *et al*, *Nano Lett.*, 2017, 17 (12), pp 7528-7532.
- [4] José M Soler *et al* 2002 *J. Phys.: Condens. Matter* 14 2745.

Figures




Figure 1: Bilayer supercell for TMDC/graphene heterostructure.




Figure 2: Electronic Band structure for WSe2 and WS2 with a vacancy in the TMDC.