

Disorder in twisted multilayer graphene: Quasicrystals and Superperiodicities

Pedro Alcázar Guerrero¹

Viet-Hung Nguyen³, Aron W. Cummings¹, Jean-Christophe Charlier³, Stephan Roche^{1,4}

¹*Catalan Institute of Nanoscience and Nanotechnology (ICN2), CSIC & BIST, Campus UAB, Bellaterra, 08193 Barcelona, Spain*

²*Department of Physics, Campus UAB, Bellaterra, 08193 Barcelona, Spain*

³*Institute of Condensed Matter and Nanosciences, Université catholique de Louvain (UCLouvain), Chemin des étoiles 8, B-1348 Louvain-La-Neuve, Belgium*

⁴*ICREA-Institució Catalana de Recerca i Estudis Avançats, 08010 Barcelona, Spain*

pedro.alcazar@icn2.cat

The stacking of 2D materials can dramatically modify the properties of those. One of the most spectacular cases is the new physics related with the stacking twist angle. That is the case of the twisted multilayer graphene, where highly correlated physics have been observed at a wide range of systems with different number of layers and twist-angle. Yet this exotic physics is only observed at ultra-clean samples and ultra-low temperatures. In this talk we will review the effects of disorder in twisted multilayers focusing on special angles such as the magic-angle ($\sim 1.05^\circ$) and the quasicrystalline ($\sim 30^\circ$). For that we will present exotic results attached to disorder as the disorder induced delocalization mechanism for the flat-band in magic angle twisted bilayer graphene (MATBLG) [1] and its disappearance when combined with a quasicrystal twisted bilayer graphene (QCTBLG) [2]. We will also cover the effects this disorder have in correlations and wave-function properties.

References

[1] PA Guerrero, VH Nguyen, JM Romeral, AW Cummings, JC Charlier, S Roche, arXiv:2401.08265

[2] PA Guerrero, VH Nguyen, AW Cummings, JC Charlier, S Roche, arXiv:2502.17069

Figures




Figure 1: **a)** Density of states of MATBLG for increasing disorder (solid $\rightarrow$ dashed $\rightarrow$ dotted). The left and right insets show the localization pattern and band structure of the clean system, respectively. **b)** Mean free path of electrons for increasing disorder strength (solid $\rightarrow$ dashed $\rightarrow$ dotted). An increase in disorder initially leads to an increase in the mean free path in the flat bands, with the usual trend recovered once its contribution dominates the Moiré potential. Insets show local densities of states for intermediate (left) and strong (right) disorder. **c)** Density of states of MATBLG and QCTBLG approximant compared to a trilayer composed by the two respective angles **d)** Diffusivity against time for the previously mentioned trilayer system and a MATBLG (in the inset), we can see the aforementioned delocalization mechanism spoiled.