Supersonic electron flow and hydraulic jump in bilayer graphene Johannes Geurs^{1,2*}

Yinjie Guo², Tatiana Webb², Itai Keren², Jack Farrell³, Takashi Taniguchi⁴, Kenji Watanabe⁴, James Hone⁵, Andrew Lucas³, Abhay Pasupathy^{2,6}, Cory R. Dean² ¹Columbia Nano Initiative, Columbia University, New York, USA ²Department of Physics, Columbia University, New York, USA ³Department of Physics, University of Colorado, Boulder, Colorado, USA

⁴Research Center for Functional Materials, National Institute for Materials Science, Tsukuba, Japan

⁵Department of Mechanical Engineering, Columbia University, New York, USA ⁶Condensed Matter Physics and Materials Science Division, Brookhaven National Lab, New York, USA

*Corresponding author: jg4424@columbia.edu

Electronic systems with global momentum conservation can be described as a hydrodynamic fluid. Clean material systems with strong carrier-carrier interactions, such as graphene [1], PdCoO₂ [2] or WTe₂ [3], can reach this regime, leading to novel electronic phenomena such as whirlpools, superballistic conductance and Poiseuille flow, all analogues of incompressible flow.

Compressible flow, where the drift velocity of the carriers is comparable to the sound velocity of the fluid and the fluid density is no longer constant, has been unexplored in electronic systems. In this work, we use the low electronic sound velocity in bilayer graphene to realize an electronic de Laval nozzle [4]. The de Laval nozzle geometry accelerates the carriers to supersonic speeds, which then relax abruptly to subsonic velocities at a shock.

Our work investigates discontinuities in electronic transport consistent with supersonic flow. Kelvin probe force measurements identify regions with supersonic flow and observe a hydraulic jump in the local potential, the equivalent of a shock wave for liquids. This is the first demonstration of compressible electron flow, and we will discuss its effects on dissipation. compressibility and reversibility of electron flow.

References

- [1] A. Lucas, K.C. Fong, J. Phys. Condens. Matter 30, 053001 (2018)
- [2] P. Moll et al. Science 351 6277, 1061-1064 (2016)
- [3] A. Aharon-Steinberg et al. Nature 607, 74-80 (2022)
- [4] K. Moors, O. Kashuba, T. L. Schmidt. arxiv:1905.01247 (2019)