

Extraordinarily transparent compact metallic metamaterials

Vincenzo Giannini

Samuel J. Palmer¹, Xiaofei Xiao¹, Nicolas Pazos-Perez², Luca Guerrini², Miguel A. Correa-Duarte³, Stefan A. Maier⁴, Richard V. Craster⁵, Ramon A. Alvarez-Puebla^{2,6}, and Vincenzo Giannini⁷

¹Department of Physics, Imperial College London, London, UK

²Department of Physical Chemistry and EMaS, Universitat Rovira i Virgili, Spain

³Department of Physical Chemistry, Singular Center for Biomedical Research (CINBIO), Southern Galicia Institute of Health Research (IISGS)

⁴Nanoinstitut München, Faculty of Physics, Ludwig-Maximilians-Universität München, Germany

⁵Department of Mathematics, Imperial College London, London, UK

⁶CREA, Passeig Lluís Companys 23, 08010 Barcelona, Spain

⁷Instituto de Estructura de la Materia (IEM-CSIC), Consejo Superior de Investigaciones Científicas, Madrid, Spain

v.giannini@csic.es

Metals are highly opaque, yet we show numerically and experimentally that densely packed arrays of metallic nanoparticles can be more transparent to infrared radiation than dielectrics such as germanium, even for arrays that are over 75% metal by volume. Despite strong interactions between the metallic particles, these arrays form effective dielectrics that are virtually dispersion-free, making possible the design of optical components that are achromatic over ultra-broadband ranges of wavelengths from a few microns up to millimetres or more.


Furthermore, the local refractive indices may be tuned by altering the size, shape, and spacing of the nanoparticles, allowing the

design of gradient-index lenses that guide and focus light on the microscale (see figure 1). The electric field is also strongly concentrated in the gaps between the metallic nanoparticles, and the simultaneous focusing and squeezing of the electric field produces strong 'doubly-enhanced' hotspots (see figure 2)

which could boost measurements made using infrared spectroscopy and other non-linear processes over a broad range of frequencies, with minimal heat production.

References

- [1] Palmer, et al. Nat. Comm. 10, 2118 (2019).



Figures

Figure 1: Schematic of a 'concentrator' gradient-index lens composed of gold nanocylinders on a triangular lattice with 50nm site-to-site separation




Figure 2: Magnetic near-fields and broadband 'doubly-enhanced' electric field hotspots
