
Unravelling the crosstalk between MXenes and extracellular vesicles

Maha Alhosani^{1*}, Olla Elkhatib^{1*}, Nabila Yasmeen^{2*}, Benjamin Chacon⁵, Jawaher Alhammadi², Laura Fusco^{3,5}, Linda Giro³, Marco Orecchioni⁴, Yury Gogotsi⁵, Yarjan Abdul Samad⁶, Anna-Maria Pappa^{2*}, and Lucia Gemma Delogu^{1,3*}

* Those authors contributed equally to this work

¹ Department of Biological Sciences, Khalifa University of Science and Technology, Abu Dhabi, UAE.

² Department of Biomedical Engineering, Khalifa University of Science and Technology, Abu Dhabi, UAE

³ Department of Biomedical Science, University of Padua, Padua, Italy

⁴ Immunology Center of Georgia, Augusta University, Augusta, Georgia

⁵ A. J. Drexel Nanomaterials Institute and Department of Materials Science and Engineering, Drexel University, Philadelphia, PA, United States.

⁶ Department of Aerospace Engineering, Khalifa University of Science Technology and Research, Abu Dhabi, UAE

lucia.delogu@ku.ac.ae

Transition metal carbides, nitrides, and carbonitrides (MXene) are emerging 2D materials with versatile chemistry, garnering significant interest in diagnostics and biomedicine [1]. Recent studies have revealed safe interactions of MXene at tissue and cellular level [2], signalling the potential possibilities for biomedical applications, such as encapsulation within extracellular vesicles (EVs) for future therapeutics. EVs are nanosized, membrane-bound particles that transfer biomolecules such as proteins, lipids, RNAs, and DNA into target cells [3]. Accumulating evidence shows that EVs have a role in intracellular communication, influencing processes like immune response and cancer progression [3,4]. Understanding the chemistry and the intracellular communication between MXene and EVs may open new avenues to revolutionize targeted drug delivery. In this study, MXene-encapsulated EVs are extracted from immune cells of THP-1 cells by mechanical extrusion protocol. The obtained changes in hydrodynamic diameter and surface charges shed light on the dynamics of the interaction of MXene with EVs at different incubation time points. Additionally, flow cytometric analysis is performed to gain deeper insight for the presence of specific extracellular and intracellular EVs surface markers. Future studies aim to delve into the potential use of MXene encapsulated EVs, extracted from THP-1 immune cells, for transfection of biomolecules and therapeutic tagging for biomedical purposes.

References

- [1] Chen, L., Dai, X., Feng, W., & Chen, Y. (2022). Biomedical applications of mxenes: From nanomedicine to Biomaterials. *Accounts of Materials Research*, 3(8), 785–798.
 - [2] Fusco, L., Gazzi, A., Shuck, C. E., Orecchioni, M., Alberti, D., D'Almeida, S. M., Rinchai, D., Ahmed, E., Elhanani, O., Rauner, M., Zavan, B., Grivel, J., Keren, L., Pasqual, G., Bedognetti, D., Ley, K., Gogotsi, Y., & Delogu, L. G. (2022). Immune profiling and multiplexed Label-Free detection of 2D MXEnes by mass cytometry and High-Dimensional imaging. *Advanced Materials*, 34(45).
 - [3] Xu, R., Greening, D. W., Zhu, H., Takahashi, N., & Simpson, R. J. (2016). Extracellular vesicle isolation and characterization: toward clinical application. *Journal of Clinical Investigation*, 126(4), 1152–1162.
 - [4] Balaj, L., Lessard, R., Dai, L., Cho, Y., Pomeroy, S. L., Breakefield, X. O., & Skog, J. (2011). Tumour microvesicles contain retrotransposon elements and amplified oncogene sequences. *Nature Communications*, 2(1).
-