

Polypeptide-based Nanoconjugates as Versatile Therapeutics

María J. Vicent

Polymer Therapeutics Lab., Prince Felipe Research Center, Valencia, Spain
mjvicent@cipf.es

Polypeptides are already playing a major role on a number of different relevant areas such as nanomedicine [1]. The physico-chemical parameters of a polypeptide-conjugate, and hence its biological performance, are defined by an intricate interplay of multiple structural factors. This highlights the need for detailed structure-activity relationship studies to develop the hierarchical strategies of polypeptide conjugate design. However, structural complexity also represents a unique opportunity, since small changes at the structural level might endow nanomedicines with outstanding and unexpected biological performance [1].

In our group, we have overcome the main classical limitations for the synthesis of defined polypeptides using precise controlled reactions followed by an adequate characterization yielding to well-defined polypeptidic architectures by NCA polymerization techniques [2]. In addition, post-polymerization techniques allow us the introduction of a variety of functionalities yielding a set of orthogonal reactive attachment sides [1,3]. Using these techniques and following a bottom-up strategy we have been able to obtain star-based polypeptide architectures with the capacity to self-assemble yielding supramolecular nanostructures with interesting properties [4]. This strategy together with an adequate polymer-drug linker design [5] enabled *in vitro* and *in vivo* evaluation, revealing a lack of toxicity, an enhanced *in vitro* cell internalization rate and significantly greater terminal and accumulation half-life *in vivo* together with a significant lymph node accumulation [4]. These results allow us to envisage these systems as promising nanocarriers for therapeutic or diagnostic applications, especially in anti-cancer treatments including lymph node metastasis and cancer immunotherapy. Proof of Concept for metastatic breast cancer [5] and for immunotherapy design in melanoma will be also shown as well as the use of this self-assembled architectures in applications such as neurodegenerative disorders [6], spinal cord injury [7] or acute kidney injury [8].

Acknowledgements

This work has been supported by European Research Council (grant ERC-CoG-2014-648831 "MyNano", Grant ERC-PoC-2018-825798 Polymune), the Spanish Ministry of Science and Innovation (SAF2016-80427-R, PID2019-108806RB-I00), Fundació Marató TV3, Fundació Step by Step and Agencia Valenciana de Innovación (AVI) INNVAL10/19/047. Part of the equipment employed in this work has been funded by Generalitat Valenciana and co-financed with FEDER funds (PO FEDER of CV 2014–2020).

References

- [1] a) Duro-Castano A. et al. (2014) *Polymers* 6, 515-551; b) Duro-Castaño A. et al. (2015) *Biomater. Sci.* 3, 1321-1334; c) Melnyk T et al. (2020) *Adv Drug Deliv Rev* 160: 136-169; d) Zagorodko O. et al. (2017) *Macromolecular Bioscience*, 17, 1600316-n/a.
- [2] a) Duro-Castano A. et al. (2015) *Mol. Pharm.* 12, 3639-3649; b) Conejos-Sánchez I. et al. (2013) *Polym. Chem.* 4, 3182-3186.
- [3] Barz M., et al. (2013) *Polym. Chemistry*, 4, 2989-2994
- [4] a) Duro-Castaño A. et al. (2017) *Adv. Mat.* 29(39): p. 1702888; b) Zagorodko O. et al. (2021) *Polym.Chem.* d1py00304f.
- [5] a) Arroyo-Crespo, J.J. et al. (2018) *Biomaterials*, 186, 8-21; b) Arroyo-Crespo, J.J., et al. (2018) *Adv. Func. Mat.* 28(22): p. 1800931
- [6] a) Duro-Castano A. et al. (2021) *Sci. Adv.* 7, eabf9180; b) Cuesta CM. et al. (2021) *Nanomedicine:NBM* 102376
- [7] Giraldo E. et al. (2021) *Biomaterials* 276, 121052
- [8] Córdoba-David G. et al. (2020) *Sci. Reports*, 10, 2056.