

Haiku graphene nanoribbons with tunable topology: emergence of localized magnetic moments

Daniel Sánchez-Portal^{1,2}

Rodrigo E. Menchón¹, Pedro Brandimarte¹, and Aran Garcia-Lekue^{1,3}

¹ Donostia International Physics Center (DIPC), 20018 Donostia-San Sebastián, Spain

² Centro de Física de Materiales CSIC-UPV/EHU, 20018 Donostia-San Sebastián, Spain

³ Ikerbasque, Basque Foundation for Science, 48013 Bilbao, Spain

daniel.sanchez@ehu.es

Abstract

Recent advances on surface-assisted synthesis open the door to engineering topological phases in atomically precise graphene nanoribbons (GNRs). However, to fully exploit their potential, a rational design is needed to achieve GNRs with optimal properties for spintronics or quantum computing applications.

Here we explore a novel family of armchair GNRs, which we name haiku-AGNRs, consisting of 5- and 7-atom wide segments. Based on ab initio simulations, we predict a tunable topological phase dependent on the density of the 7-atom wide segments, with the concomitant emergence or quenching of topological end and interface states [1]. Moreover, we derive a generalized Su-Schrieffer-Heeger (SSH) model that allows to treat haiku-AGNRs of technologically relevant lengths, thus providing valuable information for the devise of future experiments. Finally, we also present some results for B-doped periodic haiku-AGNRs in comparison with experiments performed at Prof. Pascual's lab in nanoGUNE, San Sebastian (Spain) [2].

We acknowledge financial support from Grant PID2019-107338RB-C66 funded by MCIN/AEI/10.13039/501100011033, the European Union (EU) H2020 program through the FET Open project SPRING (Grant Agreement No. 863098) and the IKUR Strategy under the collaboration agreement between Ikerbasque Foundation and DIPC

on behalf of the Dep. of Education of the Basque Government.

References

- [1] R. E. Menchón, P. Brandimarte, D. Sánchez-Portal, and A. Garcia-Lekue, submitted, (2023)
- [2] N. Friedrich, R. E. Menchón, I. Pozo, J. Hieulle, A. Vegliante, J. Li, D. Sánchez-Portal, D. Peña, A. Garcia-Lekue and J. I. Pascual, ACS Nano, 16 (2022) 14819-14826

Figures




Figure 1: DFT-SIESTA band gap of 7-(5-AGNR)_{nuc} as a function of the density ($1/n_{uc}$) of 7 widenings for periodic haiku ribbons with supercells containing n_{uc} 5-AGNR unit cells. The band gap closes separating the topological and the trivial phases. The inset highlights the presence of a defect level localized around isolated 7-widenings for very dilute systems. Notice that the labels 575-AGNR and 7-(5-AGNR)₃ denote the same system.




Figure 2: DFT-SIESTA calculation of the spin polarized end states appearing at the boundaries of the 5-AGNR portion of a mixed 5-AGNR/575-AGNR long ribbon.